



Riedgletscher VS  
Switzerland

# *Local Anesthesia for Surgical Abortions*

FIAPAC

Oct 2008, Berlin



André Seidenberg

[www.seidenberg.ch](http://www.seidenberg.ch)

# *Local Anesthesia for Surgical Abortions*

Deaths by induced abortion at  $\leq 12$  weeks' gestation

Peterson et.al. 1981

Cause of † and type of anesth.,	No. of †	No. of procedures in 100'000	Death-to-case-rate	Death-to-case-rate (95% CL)
Anesthesia-related 1972-1977	15	5'100'000	0.29	0.16 - 0.48
Local A	5	3'370'000	0.15	0.05 - 0.35
General A	10	1'730'000	0.58	0.28 - 1.06
Nonanesthesia-related, 1974-1977	24	3'980'000	0.60	0.39 - 0.90
Local A	9	2'660'000	0.34	0.15 - 0.64
General A	15	1'320'000	1.14	0.64 - 1.87

# *Local Anesthesia for Surgical Abortions*

Deaths by induced abortion at  $\leq 12$  weeks' gestation

Peterson et.al. 1981

Cause of † and type of anesth.,	No. of †	No. of procedures in 100'000	Death-to-case-rate	Death-to-case-rate (95% CL)
Anesthesia-related				
1972-1977	15	5'100'000	0.29	0.16 - 0.48
Local A	5	3'370'000	0.15	0.05 - 0.35
General A	10	1'730'000	0.58	0.28 - 1.06
Nonanesthesia-related,				
1974-1977	24	3'980'000	0.60	0.39 - 0.90
Local A	9	2'660'000	0.34	0.15 - 0.64
General A	15	1'320'000	1.14	0.64 - 1.87

# *Local Anesthesia for Surgical Abortions*

Deaths by induced abortion at  $\leq 12$  weeks' gestation

Peterson et.al. 1981

Cause of † and type of anesth.,	No. of †	No. of procedures in 100'000	Death-to-case-rate	Death-to-case-rate (95% CL)
Anesthesia-related				
1972-1977	15	5'100'000	0.29	0.16 - 0.48
Local A	5	3'370'000	0.15	0.05 - 0.35
General A	10	1'730'000	0.58	0.28 - 1.06
Nonanesthesia-related,				
1974-1977	24	3'980'000	0.60	0.39 - 0.90
Local A	9	2'660'000	0.34	0.15 - 0.64
General A	15	1'320'000	1.14	0.64 - 1.87

# *Local Anesthesia for Surgical Abortions*

Osborn et.al. 1990: General Anesthesia, a Risk Factor for Complication following Induced Abortion?

N= 10'000

Komplikationen 199 2%

Odds Ratio für Localanästhesie vs Allgemeinnarkose

Blutung 4.6 95 % CI, 2.2-9.5


Perforation 1.3 95 % CI, 0.78-2.2

Andere Komplikationen 1.6 95 % CI, 1.0-2.6

Alle Komplikationen 1.8 95 % CI, 1.4-2.5

# *Local Anesthesia for Surgical Abortions*

Schmerz von 0-10



- Menstruationsschmerzen
- Schmerz durch Infusion
- Schmerz durch Speculum
- Schmerz durch Infiltration
- Schmerz durch Dilatation
- Schmerz nach Operation
- Schmerz nach 1 Stunde
- Schmerz beim Ausgang

Aus der Dissertation von Nathalie Ambassa Mballa 2007


# *Local Anesthesia for Surgical Abortions*



*Spritze  
mit  
zwei  
Finger-  
ringen*

*Sanitex*


# *Local Anesthesia for Surgical Abortions*



*Paracervicale  
Infiltration*

*Paracervicalblock*

# *Local Anesthesia for Surgical Abortions*



*Paracervicale  
Infiltration*

*Paracervicalblock*



# *Local Anesthesia for Surgical Abortions*



*subepitheliale  
Infiltrationen*


*modifizierter  
Paracervicalblock*

# *Local Anesthesia for Surgical Abortions*



*Intracervicale  
Infiltration*

# *Local Anesthesia for Surgical Abortions*



*Transcanaliculäre  
Infiltration*